

AN ELLIPTIC EQUATION WITH NO MONOTONICITY CONDITION ON THE NONLINEARITY

GREGORY S. SPRADLIN

ABSTRACT. An elliptic PDE is studied which is a perturbation of an autonomous equation. The existence of a nontrivial solution is proven via variational methods. The domain of the equation is unbounded, which imposes a lack of compactness on the variational problem. In addition, a popular monotonicity condition on the nonlinearity is not assumed. In an earlier paper with this assumption, a solution was obtained using a simple application of topological (Brouwer) degree. Here, a more subtle degree theory argument must be used.

1. INTRODUCTION

In this paper we consider an elliptic equation of the form

$$-\Delta u + u = f(x, u), \quad x \in \mathbb{R}^N, \quad (1.1)$$

where f is a “superlinear” function of u . For large $|x|$, the equation resembles an autonomous equation

$$-\Delta u + u = f_0(u), \quad x \in \mathbb{R}^N \quad (1.2)$$

Under weak assumptions on f and f_0 , we prove the existence of a nontrivial solution u of (1.1) with $|u(x)| \rightarrow 0$ as $|x| \rightarrow \infty$.

Let $N \in \mathbb{N}^+$ and let f_0 satisfy

$$(f_1^0) \quad f_0 \in C^2(\mathbb{R}, \mathbb{R})$$

$$(f_2^0) \quad f_0(0) = 0 = f_0'(0),$$

$$(f_3^0) \quad \text{If } N > 2, \text{ there exist } a_1, a_2 > 0, s \in (1, (N+2)/(N-2)) \text{ with } |f_0'(q)| \leq a_1 + a_2|q|^{s-1} \text{ for all } q \in \mathbb{R}. \text{ If } N = 2, \text{ there exist } a_1 > 0 \text{ and a function } \varphi : \mathbb{R}^+ \rightarrow \mathbb{R} \text{ with } |f_0'(q)| \leq a_1 \exp(\varphi(|q|)) \text{ for all } q \in \mathbb{R} \text{ and } \varphi(t)/t^2 \rightarrow 0 \text{ as } t \rightarrow \infty.$$

$$(f_4^0) \quad \text{There exists } \mu > 2 \text{ such that}$$

$$0 < \mu F_0(q) \equiv \mu \int_0^q f_0(s) ds \leq f_0(q)q \quad (1.3)$$

for all $q \in \mathbb{R}$.

Let f satisfy

$$(f_1) \quad f \in C^2(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$$

$$(f_2) \quad f(x, 0) = 0 = f_q(x, 0) \text{ for all } x \in \mathbb{R}^N, \text{ where } f \equiv f(x, q)$$

2000 *Mathematics Subject Classification.* 35J20, 35J60.

Key words and phrases. Mountain Pass Theorem, variational methods, Nehari manifold, Brouwer degree, concentration-compactness.

- (f₃) If $N > 2$, there exist $a_1, a_2 > 0$, $s \in (1, (N+2)/(N-2))$ with $|f_q(x, q)| \leq a_1 + a_2|q|^{s-1}$ for all $q \in \mathbb{R}$, $x \in \mathbb{R}^N$. If $N = 2$, there exist $a_1 > 0$ and a function $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}$ with $|f_q(x, q)| \leq a_1 \exp(\varphi(|q|))$ for all $q \in \mathbb{R}$, $x \in \mathbb{R}^N$ and $\varphi(t)/t^2 \rightarrow 0$ as $t \rightarrow \infty$.
- (f₄) There exists $\mu > 2$ such that

$$0 < \mu F(x, q) \equiv \mu \int_0^q f(x, s) ds \leq f(x, q)q \quad (1.4)$$

for all $q \in \mathbb{R}$, $x \in \mathbb{R}^N$.

- (f₅) $(F(x, q) - F_0(q))/F_0(q) \rightarrow 0$ as $|x| \rightarrow \infty$, uniformly in $q \in \mathbb{R}^N \setminus \{0\}$.

In order to state the theorem, we need to outline the variational framework of the problem. Define functionals $I_0, I \in C^2(W^{1,2}(\mathbb{R}^N, \mathbb{R}), \mathbb{R})$ by

$$I_0(u) = \frac{1}{2}\|u\|^2 - \int_{\mathbb{R}^N} F_0(u(x)) dx, \quad (1.5)$$

$$I(u) = \frac{1}{2}\|u\|^2 - \int_{\mathbb{R}^N} F(x, u(x)) dx, \quad (1.6)$$

where $\|u\|$ is the standard norm on $W^{1,2}(\mathbb{R}^N, \mathbb{R})$ given by

$$\|u\|^2 = \int_{\mathbb{R}^N} |\nabla u(x)|^2 + u(x)^2 dx. \quad (1.7)$$

Critical points of I_0 correspond exactly to solutions u of (1.2) with $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$, and critical points of I correspond exactly to solutions u of (1.1) with $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

By (f₄⁰)/(f₄), F_0 and F are “superquadratic” functions of q , with $F_0(q)/q^2 \rightarrow 0$ as $q \rightarrow 0$ and $F_0(q)/q^2 \rightarrow \infty$ as $|q| \rightarrow \infty$ and $F(x, q)/q^2 \rightarrow 0$ as $q \rightarrow 0$ and $F(x, q)/q^2 \rightarrow \infty$ as $|q| \rightarrow \infty$ for all $x \in \mathbb{R}^N$, uniformly in x . Therefore $I(0) = I_0(0) = 0$, and there exists $r_0 > 0$ with $I(u) \geq \|u\|^2/3$ and $I_0(u) \geq \|u\|^2/3$ for all $u \in W^{1,2}(\mathbb{R}^N)$ with $\|u\| \leq r_0$, and there also exist $u, u_0 \in W^{1,2}(\mathbb{R}^N, \mathbb{R})$ with $I_0(u_0) < 0$ and $I(u) < 0$. So the sets of “mountain pass curves” for I_0 and I ,

$$\Gamma_0 = \{\gamma \in C([0, 1], W^{1,2}(\mathbb{R}^N, \mathbb{R})) \mid \gamma(0) = 0, I_0(\gamma(1)) < 0\}, \quad (1.8)$$

$$\Gamma = \{\gamma \in C([0, 1], W^{1,2}(\mathbb{R}^N, \mathbb{R})) \mid \gamma(0) = 0, I(\gamma(1)) < 0\}, \quad (1.9)$$

are nonempty, and the mountain-pass values

$$c_0 = \inf_{\gamma \in \Gamma_0} \max_{\theta \in [0, 1]} I_0(\gamma(\theta)) \quad (1.10)$$

$$c = \inf_{\gamma \in \Gamma} \max_{\theta \in [0, 1]} I(\gamma(\theta)) \quad (1.11)$$

are positive.

We are now in a position to state the Theorem.

Theorem 1.1. *If f_0 and f satisfy (f₁⁰)-(f₄⁰) and (f₁)-(f₅), and if there exists $\alpha > 0$ such that*

$$I_0 \text{ has no critical values in the interval } (c_0, c_0 + \alpha) \quad (1.12)$$

then there exists $\epsilon_0 = \epsilon_0(f_0) > 0$ with the following property: If f satisfies

$$|F(x, q) - F_0(q)| < \epsilon_0 F_0(q) \quad (1.13)$$

for all $x \in \mathbb{R}^N$, $q \in \mathbb{R}$, then (1.2) has a nontrivial solution $u \neq 0$ with $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

As shown in [9], (1.13) holds in a wide variety of situations.

The missing monotonicity assumption

One interesting aspect of Theorem 1.1 is a condition that is *not* assumed. We do not assume

$$\begin{aligned} &\text{For all } q \in \mathbb{R} \text{ and } x \in \mathbb{R}^N, F_0(q)/q^2 \text{ is} \\ &\quad \text{a nondecreasing function of } q \text{ for } q > 0, \\ &F_0(q)/q^2 \text{ is a nonincreasing function of } q \text{ for } q < 0, \\ &F(x, q)/q^2 \text{ is a nondecreasing function of } q \text{ for } q > 0, \text{ or} \\ &F(x, q)/q^2 \text{ is a nonincreasing function of } q \text{ for } q < 0. \end{aligned} \tag{1.14}$$

This condition holds in the power case, $F_0(q) = |q|^\alpha/\alpha$, $\alpha > 2$. The condition is due to Nehari.

If (1.14) were case, then for any $u \in W^{1,2}(\mathbb{R}^N, \mathbb{R}) \setminus \{0\}$, the mapping $s \mapsto I(su)$ would begin at 0 at $s = 0$, increase to a positive maximum, then decrease to $-\infty$ as $s \rightarrow \infty$. Defining

$$\mathcal{S} = \{u \in W^{1,2}(\mathbb{R}^N, \mathbb{R}) \setminus \{0\} \mid I'(u)u = 0\}, \tag{1.15}$$

\mathcal{S} would be a codimension-one submanifold of E , homeomorphic to the unit sphere in $W^{1,2}(\mathbb{R}^N, \mathbb{R})$ via radial projection. Any ray of the form $\{su \mid s > 0\}$ ($u \neq 0$) intersects \mathcal{S} exactly once. All nonzero critical points of I are on \mathcal{S} . Conversely, under suitable smoothness assumptions on F , any critical point of I constrained to \mathcal{S} would be a critical point of I (in the large) (see [16]). Therefore, one could work with \mathcal{S} instead of $W^{1,2}(\mathbb{R}^N, \mathbb{R})$, and look for, say, a local minimum of I constrained to \mathcal{S} (which may be easier than looking for a saddle point of I). There is another way to use (1.14): for any $u \in \mathcal{S}$, the ray from 0 passing through u can be used (after rescaling in θ) as a mountain-pass curve along which the maximum value of I is $I(u)$. Conversely, any mountain-pass curve $\gamma \in \Gamma$ intersects \mathcal{S} at least once ([6]). Therefore, one may work with points on \mathcal{S} instead of paths in Γ .

In the paper [16], a result similar to Theorem 1.1 was proven for the $N = 1$ (ODE) case. The proof of Theorem 1.1 is similar except that a simple connectivity argument must be replaced by a degree theory argument. [17] proves a version of Theorem 1.1 under the assumption (1.14). Without 1.14, the manifold \mathcal{S} must be replaced by a set with similar properties.

Define $B_1(0) = \{x \in \mathbb{R}^N \mid |x| < 1\}$, and $\overline{\Omega}$ and $\partial\Omega$ to be, respectively, the topological closure and topological boundary of Ω . It is a simple consequence of the Brouwer degree ([7]) that for any continuous function $h : \overline{B_1(0)} \rightarrow \mathbb{R}^N$ with $h(x) = x$ for all $x \in \partial B_1(0)$, there exists $x \in B_1(0)$ with $h(x) = 0$. We will need the following generalization:

Lemma 1.2. *Let $h \in C(\overline{B_1(0)} \times [0, 1], \mathbb{R}^N)$ with, for all $x \in \overline{B_1(0)}$ and $t \in [0, 1]$,*

- (i) $h(x, 0) = x = h(x, 1)$.
- (ii) $x \in \partial B_1(0) \Rightarrow h(x, t) = x$.

Then there exists a connected subset $C_0 \subset \overline{B_1(0)} \times [0, 1]$ with $(0, 0), (0, 1) \in C_0$ and $h(x, t) = 0$ for all $(x, t) \in C_0$.

Using the Brouwer degree, it is clear that under the hypotheses of Lemma 1.2, for each “horizontal slice” $\overline{B_1(0)} \times \{t\}$ of the cylinder $\overline{B_1(0)} \times [0, 1]$, there exists $x \in B_1(0)$ with $h(x, t) = 0$. The conclusion of Lemma 1.2 does not follow from this observation. It is likely that Lemma 1.2 is known; however, the author has been unable to find a proof so one is given here.

This paper is organized as follows: Section 2 contains the proof of Theorem 1.1. The proof of Lemma 1.2 is deferred until Section 3.

2. PROOF OF THEOREM 1.1

It is fairly easy to show that

$$c \leq c_0, \tag{2.1}$$

where c and c_0 are from (1.10)-(1.11): it is proven in [11] that there exists $\gamma_1 \in \Gamma_0$ with $\min_{\theta \in [0, 1]} I_0(\gamma_1(\theta)) = c_0$. Define the translation operator τ as follows: for a function u on \mathbb{R}^N and $a \in \mathbb{R}^N$, define let $\tau_a u$ be u shifted by a , that is, $(\tau_a u)(x) = u(x - a)$. Let $\epsilon > 0$. Let $\mathbf{e}_1 = \langle 1, 0, 0, \dots, 0 \rangle \in \mathbb{R}^N$ and define $\tau_{R\mathbf{e}_1} \gamma_1$ by $(\tau_{R\mathbf{e}_1} \gamma_1)(\theta) = \tau_{R\mathbf{e}_1}(\gamma_1(\theta))$. Then for large $R > 0$, by (f₅), $\tau_{R\mathbf{e}_1} \gamma_1 \in \Gamma$ and $\max_{\theta \in [0, 1]} I((\tau_{R\mathbf{e}_1} \gamma_1)(\theta)) < c_0 + \epsilon$. Since $\epsilon > 0$ was arbitrary, $c \leq c_0$.

A Palais-Smale sequence for I is a sequence $(u_m) \subset W^{1,2}(\mathbb{R}^N, \mathbb{R})$ with $(I(u_m))$ convergent and $\|I'(u_m)\| \rightarrow 0$ as $m \rightarrow \infty$. It is well-known that I fails the “Palais-Smale condition.” That is, a Palais-Smale sequence need not converge. However, the following proposition states that a Palais-Smale sequence “splits” into the sum of a critical point of I and translates of critical points of I_0 :

Proposition 2.1. *If $(u_m) \subset W^{1,2}(\mathbb{R}^N, \mathbb{R})$ with $I'(u_m) \rightarrow 0$ and $I(u_m) \rightarrow a > 0$, then there exist $k \geq 0$, $v_0, v_1, \dots, v_k \in W^{1,2}(\mathbb{R}^N, \mathbb{R})$, and sequences $(x_m^i)_{m \geq 1}^{1 \leq i \leq k} \subset \mathbb{R}^N$, such that*

- (i) $I'(v_0) = 0$
- (ii) $I'_0(v_i) = 0$ for all $i = 1, \dots, k$

and along a subsequence (also denoted (u_m))

- (iii) $\|u_m - (v_0 + \sum_{i=1}^k \tau_{x_m^i} v_i)\| \rightarrow 0$ as $m \rightarrow \infty$
- (iv) $|x_m^i| \rightarrow \infty$ as $m \rightarrow \infty$ for $i = 1, \dots, k$
- (v) $|x_m^i - x_m^j| \rightarrow \infty$ as $m \rightarrow \infty$ for all $i \neq j$
- (iii) $I(v_0) + \sum_{i=1}^k I_0(v_i) = a$

A proof for the case of x -periodic F is found in [6], and essentially the same proof works here. Similar propositions for nonperiodic coefficient functions, for both ODE and PDE, are found in [5], [1], and [18], for example. All are inspired by the “concentration-compactness” theorems of P. -L. Lions ([12]).

If $c < c_0$, then by standard deformation arguments ([15]), there exists a Palais-Smale sequence (u_m) with $I(u_m) \rightarrow c$. By [11], the smallest nonzero critical value of I_0 is c_0 . Applying Proposition 2.1, we obtain $k = 0$, and (u_m) has a convergent subsequence, proving Theorem 1.1. So assume from now on that

$$c = c_0. \tag{2.2}$$

For $u \in L^2(\mathbb{R}^N, \mathbb{R}) \setminus \{0\}$ and $i \in \{1, \dots, N\}$, define \mathcal{L}_i , the i th component of the “location” of u , by

$$\int_{\mathbb{R}^N} u^2 \tan^{-1}(x_i - \mathcal{L}_i(u)) dx = 0 \quad (2.3)$$

and the “location” of u by

$$\mathcal{L}(u) = (\mathcal{L}_1(u), \dots, \mathcal{L}_N(u)) \in \mathbb{R}^N. \quad (2.4)$$

We are ready to begin the minimax argument. First we construct a mountain-pass curve γ_0 with some special properties:

Lemma 2.2. *There exists $\gamma_0 \in \Gamma_0$ such that for all $\theta \in [0, 1]$,*

- (i) $I_0(\gamma_0(\theta)) \leq c_0$.
- (ii) $\theta > 0 \Rightarrow \gamma_0(\theta) \neq 0$.
- (iii) $\theta \leq 1/2 \Rightarrow I_0(\gamma_0(\theta)) \leq c_0/2$.
- (iv) $\theta > 0 \Rightarrow \mathcal{L}(\gamma_0(\theta)) = 0$.

Proof: by [10], there exists $\gamma_1 \in \Gamma_0$ with $\max_{\theta \in [0, 1]} I_0(\gamma_1(\theta)) = c_0$. Assume without loss of generality that $\gamma_1(\theta) \neq 0$ for $\theta > 0$. By rescaling in θ if necessary, assume that $I_0(\gamma_1(\theta)) \leq c_0/2$ for $\theta \leq 1/2$. Finally, define γ_0 by $\gamma_0(0) = 0$, $\gamma_0(\theta) = \tau_{-\mathcal{L}(\gamma_1(\theta))}\gamma_1(\theta)$ for $\theta > 0$.

Assume ϵ_0 in (1.13) is small enough so that for all $x \in \mathbb{R}^N$ and $\theta \in [0, 1]$,

$$I(\tau_x(\gamma_0(\theta))) < \min(2c_0, c_0 + \alpha) \text{ and } I(\tau_x(\gamma_0(1))) < 0, \quad (2.5)$$

where α is from (1.12).

A substitute for \mathcal{S}

Using the mountain-pass geometry of I and the fact that Palais-Smale sequences of I are bounded in norm ([6]), we construct a set which has similar properties to \mathcal{S} , described in Section 1. Let ∇I denote the gradient of I , that is, $(\nabla I(u), w) = I'(u)w$ for all $u, w \in W^{1,2}(\mathbb{R}^N, \mathbb{R})$. Here, (\cdot, \cdot) is the usual inner product defined by $(u, w) = \int_{\mathbb{R}^N} \nabla u \cdot \nabla w + uw dx$. Let $\varphi : W^{1,2}(\mathbb{R}^N, \mathbb{R}) \rightarrow \mathbb{R}$ be locally Lipschitz, with $I(u) \geq -1 \Rightarrow \varphi(u) = 1$ and $I(u) \leq -2 \Rightarrow \varphi(u) = 0$. Let η be the solution of the initial value problem

$$\frac{d\eta}{ds} = -\varphi(\eta)\nabla I(u), \quad \eta(0, u) = u. \quad (2.6)$$

In [18] it is proven that η is well-defined on $\mathbb{R}^+ \times W^{1,2}(\mathbb{R}^N)$. Let \mathcal{B} be the basin of attraction of 0 under the flow η , that is,

$$\mathcal{B} = \{u \in W^{1,2}(\mathbb{R}^N, \mathbb{R}) \mid \eta(s, u) \rightarrow 0 \text{ as } s \rightarrow \infty\} \quad (2.7)$$

\mathcal{B} is an open neighborhood of 0 ([18]). Let $\partial\mathcal{B}$ be the topological boundary of \mathcal{B} in $W^{1,2}(\mathbb{R}^N, \mathbb{R})$. $\partial\mathcal{B}$ has some properties in common with \mathcal{S} . For example, for any $\gamma \in \Gamma$, $\gamma([0, 1])$ intersects $\partial\mathcal{B}$ at least once.

A pseudo-gradient vector field for I' may be used in place of ∇I , in which case \mathcal{B} and $\partial\mathcal{B}$ would be different, but the ensuing arguments would be the same.

Let

$$c^+ = \inf\{I(u) \mid u \in \partial\mathcal{B}, |\mathcal{L}(u)| \leq 1\}. \quad (2.8)$$

The reason for the label “ c^+ ” will become apparent in a moment. From now on, let us assume

$$I \text{ has no critical values in } (0, c_0] = (0, c]. \quad (2.9)$$

This will lead to the conclusion that I has a critical value greater than c_0 .

We claim that under assumptions (2.2) and (2.9),

$$c^+ > c_0. \quad (2.10)$$

To prove the claim, suppose first that $c^+ < c_0$. Then there exists $u_0 \in \partial\mathcal{B}$ with $I(u_0) < c_0$. Define $u_n = \eta(n, u_0)$. By arguments of [18] and [5], $\|I'(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$ and there exists $b \in (0, c_0)$ with $I(u_n) \rightarrow b$. By [11], I_0 has no critical values between 0 and c_0 . Therefore, Proposition 2.1, with $k = 0$, implies that (u_n) converges along a subsequence to a critical point w of I with $0 < I(w) < c_0$. This contradicts assumption (2.9).

Next, suppose that $c^+ = c_0$. Then there exists a sequence $(u_n) \subset \partial\mathcal{B}$ with $I(u_n) \rightarrow c_0$ as $n \rightarrow \infty$. By the arguments of [18], $I'(u_n) \rightarrow 0$ as $n \rightarrow \infty$; to prove, suppose otherwise. Then there exists $b > 0$ and a subsequence of (u_n) (also called (u_n)) along which $\|I'(u_n)\| > b$. Since $\partial\mathcal{B}$ is forward- η -invariant ([18]), $\eta(1, u_n) \in \partial\mathcal{B}$ for all n . Since $(\eta(1, u_n))_{n \geq 1}$ is bounded ([18]) and I' is Lipschitz on bounded subsets of $W^{1,2}(\mathbb{R}^N, \mathbb{R})$, for large n , $\eta(1, u_n) \in \partial\mathcal{B}$ with $I(\eta(1, u_n)) < c_0$. By the argument above, this implies that I has a critical value in $(0, c_0)$, contradicting assumption (2.2). Thus $I'(u_n) \rightarrow 0$ as $n \rightarrow \infty$. Applying Proposition 2.1 and using the fact that $|\mathcal{L}(u_n)| \leq 1$ for all n , (u_n) converges along a subsequence to a critical point of I , contradicting assumption (2.9). (2.10) is proven.

Let $R > 0$ be big enough so that for all $x \in \partial B_R(0) \subset \mathbb{R}^N$ and $\theta \in [0, 1]$,

$$I(\tau_x \gamma_0(\theta)) < c^+. \quad (2.11)$$

This is possible by (1.13), (2.10), and Lemma 2.2(i). Define the minimax class

$$\begin{aligned} \mathcal{H} = \{ & h \in C(\overline{B_R(0)} \times [0, 1], W^{1,2}(\mathbb{R}^N, \mathbb{R})) \mid \\ & \text{for all } x \in \overline{B_R(0)} \text{ and } t \in [0, 1], \\ & t > 0 \Rightarrow h(x, t) \neq 0 \\ & 0 \leq t \leq 1/2 \Rightarrow h(x, t) = \tau_x \gamma_0(t) \\ & x \in \partial B_R(0) \Rightarrow h(x, t) = \tau_x \gamma_0(t) \\ & h(x, 1) = \tau_x \gamma_0(1) \} \end{aligned}$$

and the minimax value

$$h_0 = \inf_{h \in \mathcal{H}} \max_{(x,t) \in \overline{B_R(0)} \times [0,1]} I(h(x, t)). \quad (2.12)$$

We claim

$$c_0 < c^+ \leq h_0 < \min(2c_0, c_0 + \alpha). \quad (2.13)$$

Proof of Claim: define $\bar{h} \in \mathcal{H}$ by $\bar{h}(x, t) = \tau_x(\gamma_0(t))$. Then $\bar{h} \in \mathcal{H}$ and by (2.5), $\max_{(x,t) \in \overline{B_R(0)} \times [0,1]} \bar{h}(x, t) < \min(2c_0, c_0 + \alpha)$. Therefore $h_0 < \min(2c_0, c_0 + \alpha)$.

Next, let $h \in \mathcal{H}$. By Lemma 1.2, and a suitable rescaling of x and t , there exists a connected set $C_2 \subset B_R(0) \times [1/2, 1]$ with $(0, 1/2), (0, 1) \in C_2$ and along which for all $(x, t) \in C_2$,

$$\mathcal{L}(h(x, t)) = 0. \quad (2.14)$$

Joining C_2 with the segment $\{0\} \times [0, 1/2]$, we obtain a connected set $C_3 \subset B_R(0) \times [0, 1]$ such that $(0, 0), (0, 1) \in C_3$ and for all $(x, t) \in C_3$, $\mathcal{L}(h(x, t)) = 0$. C_3 is not

necessarily path-connected, so let $r > 0$ be small enough so that for all

$$\begin{aligned} (x, t) \in N_r(C_3) &\equiv \{(y, s) \in B_R(0) \times [0, 1] \mid \\ &\exists (x', t') \in B_R(0) \times [0, 1] \text{ with } |y - x'|^2 + (s - t')^2 < r^2\}, \\ &|\mathcal{L}(h(x, t))| < 1. \end{aligned} \quad (2.15)$$

$N_r(C_3)$ is path-connected ([20]), so there exists a path $g \in C([0, 1], N_r(C_3))$ with $g(0) = (0, 0)$, $g(1) = (0, 1)$, and $g(\theta) \in N_r(C_3)$ for all $\theta \in [0, 1]$. If we define $\tilde{\gamma} \in \Gamma$ by $\tilde{\gamma}(\theta) = h(g(\theta))$, then $|\mathcal{L}(\tilde{\gamma}(\theta))| < 1$ for all $\theta \in [0, 1]$. Since $\tilde{\gamma}(0) = 0$ and $I(\tilde{\gamma}(1)) < 0$, there exists $\theta^* \in [0, 1]$ with $\tilde{\gamma}(\theta^*) \in \partial\mathcal{B}$. By the definition of c^+ (2.8), $I(\tilde{\gamma}(\theta^*)) \geq c^+$.

Since h was an arbitrary element of \mathcal{H} , $h_0 \geq c^+$.

By standard deformation arguments, such as described in [15], there exists a Palais-Smale sequence $(u_n) \subset W^{1,2}(\mathbb{R}^N, \mathbb{R})$ with $I'(u_n) \rightarrow 0$ and $I(u_n) \rightarrow h_0$ as $n \rightarrow \infty$. $c_0 < h_0 < \min(2c_0, c_0 + \alpha)$. Apply Proposition 2.1 to (u_n) . Since I_0 has no positive critical values smaller than c_0 ([11]), $k \leq 1$. By (2.9), (u_n) converges along a subsequence to a critical point z of I , with $I(z) = h_0$. Theorem 1.1 is proven.

3. A DEGREE-THEORETIC LEMMA

Here, we prove Lemma 1.2. Let h be as in the hypotheses of the lemma. For $l > 0$, define $\mathcal{A}_l \subset \overline{B_1(0)} \times [0, 1]$ by

$$\mathcal{A}_l = \{(x, t) \in \overline{B_1(0)} \times [0, 1] \mid |f(x, t)| < l\}. \quad (3.1)$$

\mathcal{A}_l is an open neighborhood of $(0, 0)$. Let C_l be the component of \mathcal{A}_l containing $(0, 0)$. We will prove the following claim:

$$\text{For all } \epsilon > 0, (0, 1) \in C_\epsilon. \quad (3.2)$$

Then we will use the C_ϵ 's to construct C_0 . For $l > 0$ and $t \in [0, 1]$, define

$$C_l^t = \{x \in \overline{B_1(0)} \mid (x, t) \in C_l\}. \quad (3.3)$$

Fix $\epsilon \in (0, 1)$. Define $\phi : [0, 1] \rightarrow \mathbb{Z}$ by

$$\phi(t) = d(h(\cdot, t), C_\epsilon^t, 0), \quad (3.4)$$

where d is the topological Brouwer degree ([7]). We will prove $\phi(t) = 1$ for all $t \in [0, 1]$, in particular $\phi(1) = 1$, so (3.2) is satisfied.

f is continuous on a compact domain, so f is uniformly continuous. Let $\rho > 0$ be small enough so that for all $x \in \overline{B_1(0)}$ and $t_1, t_2 \in [0, 1]$,

$$|t_1 - t_2| < \rho \Rightarrow |h(x, t_1) - h(x, t_2)| < \epsilon/4. \quad (3.5)$$

Clearly

$$\phi(0) = d(\text{id}, B_\epsilon(0), 0) = 1. \quad (3.6)$$

Let $0 \leq t_1 < t_2 \leq 1$ with $t_2 - t_1 < \rho$. We will show $\phi(t_1) = \phi(t_2)$, proving that ϕ is constant, which by (3.6), implies (3.2).

Ω is nonempty. For all $x \in \partial C_\epsilon^{t_1}$, $|h(x, t_1)| = \epsilon$, so by (3.5),

$$x \in \partial C_\epsilon^{t_1} \Rightarrow |h(x, t_1)| \geq \frac{3}{4}\epsilon. \quad (3.7)$$

By the additivity property of d ([7]),

$$\begin{aligned}\phi(t_2) &\equiv d(f(\cdot, t_2), C_\epsilon^{t_2}, 0) = \\ &= d(f(\cdot, t_2), C_\epsilon^{t_2} \setminus \overline{C_\epsilon^{t_1}}, 0) + d(f(\cdot, t_2), C_\epsilon^{t_1} \cap C_\epsilon^{t_2}, 0).\end{aligned}\tag{3.8}$$

We will show:

$$\text{There does not exist } x \in C_\epsilon^{t_2} \setminus \overline{C_\epsilon^{t_1}} \text{ with } h(x, t_2) = 0.\tag{3.9}$$

Suppose such an x exists. Then by (3.5), $|h| < \epsilon/4$ on the segment $\{x\} \times [t_1, t_2]$. $x \in C_\epsilon^{t_2}$, so $(x, t_2) \in C_\epsilon$, and by the definition of C_ϵ , $(x, t_1) \in C_\epsilon$, and $x \in C_\epsilon^{t_1}$, contradicting $x \in C_\epsilon^{t_2} \setminus \overline{C_\epsilon^{t_1}}$. So (3.9) is true. Therefore by (3.8),

$$\phi(t_2) = d(f(\cdot, t_2), C_\epsilon^{t_1} \cap C_\epsilon^{t_2}, 0).\tag{3.10}$$

By the same argument, switching the roles of t_1 and t_2 ,

$$\phi(t_1) = d(f(\cdot, t_1), C_\epsilon^{t_1} \cap C_\epsilon^{t_2}, 0).\tag{3.11}$$

For all $t \in [t_1, t_2]$ and $x \in \partial C_\epsilon^{t_1} \cup \partial C_\epsilon^{t_2}$, (3.5) gives $|h(x, t_1)| > 3\epsilon/4$ and $|h(x, t) - h(x, t_1)| < \epsilon/4$. Therefore by the homotopy invariance property of the degree ([7]),

$$\begin{aligned}\phi(t_1) &= d(f(\cdot, t_1), C_\epsilon^{t_1} \cap C_\epsilon^{t_2}, 0) = \\ &= d(f(\cdot, t_2), C_\epsilon^{t_1} \cap C_\epsilon^{t_2}, 0) = \phi(t_2).\end{aligned}\tag{3.12}$$

$\phi(0) = 1$ and $\phi(t_1) = \phi(t_2)$ for any $t_1 < t_2$ with $t_1, t_2 \in [0, 1]$ and $t_2 - t_1 < \rho$. Therefore ϕ is constant, and $\phi(1) = 1$. Therefore $(0, 1) \in C_\epsilon$.

Now let

$$C_0 = \bigcap_{\epsilon > 0} C_\epsilon.\tag{3.13}$$

Each C_ϵ is a connected set containing $(0, 0)$ and $(0, 1)$, so it is easy to show that C_0 is a connected set containing $(0, 0)$ and $(0, 1)$, and clearly for all $(x, t) \in C_0$, $h(x, t) = 0$.

REFERENCES

- [1] F. Alessio and P. Montecchiari, *Multibump solutions for a class of Lagrangian systems slowly oscillating at infinity*, Annales de l'Institut Henri Poincaré, Vol. 16 (1999), No. 1, 107-135.
- [2] A. Bahri and Y.-Y. Li, *On a Min-Max Procedure for the Existence of a Positive Solution for a Certain Scalar Field Equation in \mathbb{R}^N* , Revista Iberoamericana, Vol. 6 (1990), 1-17.
- [3] P. Caldiroli, *A New Proof of the Existence of Homoclinic Orbits for a Class of Autonomous Second Order Hamiltonian Systems in \mathbb{R}^N* , Math. Nachr., Vol. 187 (1997), 19-27.
- [4] P. Caldiroli, P. Montecchiari, *Homoclinic orbits for second order Hamiltonian systems with potential changing sign*, Comm. Appl. Nonlinear Anal., Vol. 1 (1994), No. 2, 97-129.
- [5] V. Coti Zelati, P. Montecchiari, and M. Nolasco, *Multibump solutions for a class of second order, almost periodic Hamiltonian systems*, Nonlinear Ordinary Differential Equations and Applications, Vol. 4 (1997), No. 1, 77-99.
- [6] V. Coti Zelati and P. Rabinowitz, *Homoclinic Orbits for Second Order Hamiltonian Systems Possessing Superquadratic Potentials*, Journal of the American Mathematical Society, Vol. 4 (1991), 693-627.
- [7] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, New York, 1985.
- [8] M. Estaban and P.-L. Lions, *Existence and non existence results for semilinear elliptic problems in unbounded domains*, Proc. Roy. Soc. Edinburgh, Vol. 93 (1982), 1-14.
- [9] B. Franchi, E. Lanconelli, and J. Serrin, *Existence and Uniqueness of Nonnegative Solutions of Quasilinear Equations in \mathbb{R}^N* , Advances in Mathematics, Vol. 118, (1996), 177-243.
- [10] L. Jeanjean, K. Tanaka, *A Note on a Mountain Pass Characterization of Least Energy Solutions*, Adv. Nonlinear Stud., Vol. 3, (2003), 445-455.

- [11] L. Jeanjean, K. Tanaka, *A remark on least energy solutions in \mathbb{R}^N* , Proc. Amer. Math. Soc., Vol. 131 (2003), 2399-2408.
- [12] P. L. Lions, *The concentration-compactness principle in the calculus of variations. The locally compact case.*, Annales de l'Institut Henri Poincaré, Vol. 1 (1984), 102-145 and 223-283.
- [13] J. Mawhin and M. Willem, "Critical Point Theory and Hamiltonian Systems," Springer-Verlag, New York, 1989.
- [14] P. Rabinowitz, *Homoclinic Orbits for a class of Hamiltonian Systems*, Proc. Roy. Soc. Edinburgh, Sect. A, Vol. 114 (1990), 33-38.
- [15] P. Rabinowitz, "Minimax Methods in Critical Point Theory with Applications to Differential Equations," C. B. M. S. Regional Conf. Series in Math., No. 65, Amer. Math. Soc., Providence, 1986.
- [16] G. Spradlin, *Existence of Solutions to a Hamiltonian System without Convexity Condition on the Nonlinearity*, Electronic Journal of Differential Equations, Vol. 2004 (2004), No. 21 1-13.
- [17] G. Spradlin, *A Perturbation of a Periodic Hamiltonian System*, Nonlinear Analysis, Theory, Methods, and Applications, Vol. 38, No. 8 (1999), 1003-1022.
- [18] G. Spradlin *Interacting Near-Solutions of a Hamiltonian System*, Calc. Var. PDE, Vol. 22 (2005), 447-464.
- [19] E. Serra, M. Tarallo, and S. Terracini, *On the existence of homoclinic solutions to almost periodic second order systems*, Annales de l'Institut Henri Poincaré, Vol. 13 (1996), 783-812.
- [20] G. Whyburn, *Topological Analysis*, Princeton University Press, 1964.

GREGORY S. SPRADLIN
DEPARTMENT OF MATHEMATICS, EMBRY-RIDDLE AERONAUTICAL UNIVERSITY, DAYTONA BEACH,
FLORIDA 32114-3900, USA
E-mail address: spradlig@erau.edu